

Cherry leaf roll virus Abundant on *Betula pubescens* in Finland

Risto Jalkanen, Carmen Büttner and Susanne von Bargaen

Jalkanen, R., Büttner, C. & von Bargaen, S. 2007. Cherry leaf roll virus abundant on *Betula pubescens* in Finland. *Silva Fennica* 41(4): 755–762.

Virus-related symptoms such as vein banding, leaf roll, chlorosis and subsequent necrosis on birch leaves were increasingly recorded throughout Finland since 2002. They are widespread in this country and have also been detected in northern Norway and Sweden. Symptomatic foliage has so far been found on *Betula pendula*, *B. pubescens*, *B. pubescens* subsp. *czerepanovii*, and *B. nana*.

A Cherry leaf roll virus (CLR) specific IC-RT-PCR was applied to young leaves, buds and catkins of symptomatic shoots of nineteen pubescent and one silver birch trees grown in the centre of Rovaniemi, Finland. CLR was found in seventeen *B. pubescens* trees. This is the first time that *B. pubescens* has been confirmed to be a host species for CLR in Finland. Nor has CLR been recorded earlier in northern Finland.

Keywords *Betula pendula*, *Betula nana*, *Betula pubescens* subsp. *czerepanovii*, virology

Authors' addresses Jalkanen, Finnish Forest Research Institute, Rovaniemi Research Unit, P.O. Box 16, FI-96301 Rovaniemi, Finland; Büttner and von Bargaen, Humboldt Universität zu Berlin, Institut für Gartenbauwissenschaften, Fachgebiet Phytomedizin, Lentzeallee 55/57, 14195 Berlin, Germany **E-mail** risto.jalkanen@metla.fi

Received 4 April 2007 **Revised** 6 September 2007 **Accepted** 17 September 2007

Available at <http://www.metla.fi/silvafennica/full/sf41/sf414755.pdf>

1 Background

Since the summer of 2002, increasing numbers of birch trees (*Betula* spp.) of various sizes and ages with foliar disorders of unknown origin were discovered throughout Fennoscandia. Symptomatic birches showing vein banding, leaf roll and chlorosis with subsequent necrosis were found from the southern coast of Finland to the conifer

tree-line area in northern Finland, in the sub arctic zone of northern Norway as well as in Sweden (Fig. 1). Affected birch species were silver (*Betula pendula* Roth), pubescent (*B. pubescens* Ehrh.), mountain (*B. pubescens* subsp. *czerepanovii* (Orlova) Hämet-Ahti) and dwarf birch (*B. nana* L.). Diseased birch trees were either growing alone or in groups. The phenomenon occurred in trees and bushes on mineral sites and peatland, in




Fig. 1. Locations, where *Cherry leaf roll virus* -type symptoms have been recorded on *Betula* spp. in Fennoscandia. The black square represents 25 records in Rovaniemi, N Finland.

inhabited areas such parks, and forests, and in valleys as well as exactly at the birch alpine tree line. Leaf symptoms appeared either in a small part, even a branch tip, or in most parts of the green canopy (Fig. 2). Tops of taller trees seemed to be least affected. Leaf symptoms of this type have neither been found ever to this extent in Finland nor confirmed for the causal agent on *B. pubescens* or other birch species except *B. pendula*.

2 Symptom Description

First symptoms on *B. pubescens* appeared soon after full expansion of first leaves and after the primary shoots at tips of first-order branches had grown a few centimetres at the end of May (Rovaniemi 2006). At this time, chlorotic vein banding, lighter than the pale green early summer foliage colour, developed and affected leaves started to

show leaf roll, leading to a convex appearance (Fig. 3a). In some cases chlorotic line patterns were more irregular and concentrated on the leaf tips. First symptoms appeared in the primary leaves, but not every leaf was affected. While symptom-free leaves expanded to regular size with a normal shape and green colour, the final size of the symptomatic leaves remained noticeably smaller.

In further development all leaves at the tip part of the new shoot exhibited symptoms. From July onwards the general colour of the symptomatic foliage increasingly changed from a lighter green to yellow and, in most cases, exhibited strong leaf roll symptoms. Chlorotic bands were clearly visible either between the main veins or on both sides of a vein, forming either diamond structures or chlorotic areas widening towards the leaf edge. Sometimes the whole interveinal space was chlorotic (Fig. 3c), while veins and adjacent tissues remained green most of the season. Ring spots were absent but patterns of the mosaic type occurred in some leaves of normal size (Fig. 3b). Throughout the summer strongest symptoms were present at the leaf tips, where chlorotic lesions gradually turned necrotic (Fig. 2d). Leaf roll symptoms increased in time as well (Fig. 3d, also 3a and 3c). Additionally, in the very dry summer of 2006, symptomatic leaves were drooping, thus the affected parts of the canopy resembled to suffer from drought (Fig. 2b). Further, distorted foliage, either due to hypertrophic growth of interveinal tissue or reduced vein extension was common (Fig. 2e, 3c).

In late summer the symptomatic parts of the canopies could easily be distinguished from symptom-free parts or the neighbouring crowns due to their pale colour (Fig. 4). Normally, a lot of parasitic fungi inhabited birch leaves at the end of the growing season so that fungal infections partly hid symptoms caused by other diseases. In this respect fungal pathogens infected birch foliage exceptionally scanty in the summer of 2006, this allowed the studied phenomenon to appear well.

– Shedding time of symptomatic leaves in respect to healthy ones seemed to vary depending on site, tree and even location within the canopy. Once a tree exhibited the above-described symptoms, they clearly appeared to have spread the following year. In fact, the entire canopy could become symptomatic in a few years.




Fig. 2. Virus-like symptoms found throughout Fennoscandia. **a)** On a shoot tip of *Betula pubescens* (sample tree no. 10) in the town Rovaniemi, N Finland, 19.8.2006; **b)** in the lower canopy of *B. pendula* with healthy and infected leaves/shoots in Jyväskylä, central Finland, 1.9.2006; **c)** in a shoot of *B. pubescens* var. *czerepanovii* in Neiden, NE Norway, 19.7.2006; **d)** necrotic lesions on *B. pubescens* leaves, Inari village, Inari, N Finland 22.8.2006; and **e)** hypertrophic leaf growth of *B. pubescens* tree, in the town Luleå, N Sweden, 14.9.2004.




Fig. 3. Symptoms of *Cherry leaf roll virus*, CLRV, infected *Betula pubescens*, town Rovaniemi, N Finland **a**) Leaf-roll symptoms and beginning chlorosis soon after the shoot flush, sample tree no. 8, 1.6.2006; **b**) mosaic pattern, sample tree no. 14, 21.9.2006; **c**) typical chlorotic vein banding, line pattern and leaf roll, sample tree no. 2, 14.7.2006; and **d**) leaf-roll symptoms together with line pattern, sample tree no. 2, 4.8.2006.

The syndrome of birch trees described here led to the hypothesis of a viral etiology. Although, several viruses like *Tobacco rattle virus* (TRV, Schmelzer 1972) and *Apple mosaic virus* (ApMV, Grüntzig et al. 1996) have been reported to infect *Betula* species, symptoms resembled those described for *Cherry leaf roll virus*, CLRV (Schmelzer 1972, Cooper and Atkinson 1975), known to have a wide geographic distribution (Rebenstorf et al. 2006). To confirm our suspi-

cion, we aimed to apply the IC-RT-PCR technique to test the presence of CLRV in northern Finnish birch trees.

3 Material and Methods

Nineteen pubescent birches and one silver birch, which exhibited the above-mentioned symptoms,



Fig. 4. Birch habitus with typical *Cherry leaf roll virus*, CLRV, type symptoms in the entire canopy in Rovaniemi, N Finland, 4.8.2006.

were numbered (1 to 20) and sampled in the centre of the Rovaniemi town in northern Finland (66°29'...66°31' N) in September 21, 2006. At the time of sampling only one symptomatic silver birch in Rovaniemi was known. The sampled

trees were 3 to 14 m tall, and their green canopy exhibited slight, moderate or strong symptoms. In the slightest case, symptoms appeared as a rule only in the lower parts of the green canopy. One to two twigs with leaves, buds and, if possible, catkins were sampled from symptomatic areas of the lower canopy. In case of trees nos. 18–20 symptoms were different or became already diffuse due to the autumn colouration of leaves. Sampled material was stored wrapped in moist paper towels in plastic bags at +5 °C.

The samples were analysed within one week from sampling by application of a modified CLRV specific immuno capture reverse transcriptase polymerase chain reaction (IC-RT-PCR) developed by Werner et al. (1997). One symptomatic leaf, two to three buds and one catkin, if available, per tree were tested. To apply for routine testing, the protocol was shortened by omitting denaturation steps and carried out as a single tube reaction assay as well as reducing required chemicals and enzymes, in short: One 0.2 ml reaction tube per sample was coated with 10 µl antibody solution (1:200 dilution of IgG in carbonate buffer, pH 9.6) for 2 h at 37 °C. The polyclonal antibody was raised in rabbit against a CLRV isolate E603 from a black elderberry, *Sambucus nigra* L. (in Rebenstorf et al. 2006). Immunisation and purification of IgG were done using standard protocols (Harlow and Lane 1988). After washing the tubes three times with 150 µl phosphate buffered saline containing 0.05% (v/v) Tween (PBS-Tween), 10 µl homogenised plant material (1:10 dilution in PBS-Tween supplemented with 2% (w/v) polyvinylpyrrolidone 40) was applied and incubated over night at 4 °C. Unbound material was washed away as described above and 10 µl of reverse transcriptase mix, which consists of 50 mM Tris-HCl pH 8.3, 50 mM KCl, 4 mM MgCl₂, 10 mM dithiothreitol, 1 mM each dNTP, 50 pmol first strand primer (RW1, 5' GTCGGAAAGATTACGTAAAAGG) and 200 units M-MuLV-RTase (Fermentas), was added to reaction tubes and incubated for at least 1 h at 37 °C. Afterwards 40 µl of PCR mix – 12.5 mM Tris, 62.5 mM KCl, 0.125% (v/v) TritonX100, 0.25 mM each dNTP, 2.5 mM MgCl₂, 25 pmol each first and second strand primer (RW2, 5' TGGCGACCGTGTAACGGCA) and 5 units Taq-Polymerase (Promega) – was added directly

to the cDNA reaction and subjected to PCR in a robocycler (Stratagene) using the following program: 1× 120 s 94 °C; 30× 60 s 94 °C, 45 s 55 °C, 60 s 72 °C; 1× 5 min 72 °C. PCR-products were analysed by separation in 1% agarose gels and visualised after staining with ethidiumbromide under uv-light.

In parallel, samples of six trees (nos. 2, 8, 14, 15, 17 and 20) were checked for virus particles stained with 2% (w/v) uranylacetate by transmission electron microscopy (TEM).

4 Results and Discussion

CLRV was detected in leaves, buds and/or catkins in 85% of the sampled trees (Table 1). IC-RT-PCR specifically amplified a single fragment of the expected size (412 bp), from sampled tissues. This was in accordance with amplification products received from CLRV infected woody control plants (*Sambucus nigra* and *B. pendula*), while no fragments were obtained from the water control. Only in three birches IC-RT-PCR results were negative. Symptoms of the two CLRV negative *B. pubescens* trees differed from the patterns described above and referenced in the literature for CLRV. The observed leaf-roll symptoms might be due to the very severe drought in the sampling area in 2006 (Monthly weather report 2006), together with the appearance of autumn colours. The third tree with a negative result was the only sampled *B. pendula*, but with similar symptoms as the trees, which resulted in CLRV infected. The failure to detect CLRV in this sample might be due to improper sample material, because only very few shoots were symptomatic. Although, CLRV detection is usually possible in leaves from woody hosts showing chlorotic spots and line pattern, virus concentration is often low in these plants and the virus is irregularly distributed within the canopy (Bandte and Büttner 2001). Leaves of the tested silver birch tree were already necrotising at the end of the vegetation period, which also might be a cause for lack of detectability of CLRV. Secondary plant compounds in sampled material such as phenols and polysaccharides, which are known inhibitors of PCR (Demeke and Adams 1992, MacKenzie et al.

Table 1. Presence of *Cherry leaf roll virus*, CLRV, in *Betula* spp. growing along the streets in the centre of Rovaniemi, Finland.

Tree no.	Species	Symptom appearance ^{a)}	CLRV detection ^{b)}		
			Leaves	Buds	Catkins
1	<i>B. pubescens</i>	C, T, o	+	-	-
2	<i>B. pubescens</i>	C, T, p	+	-	na
3	<i>B. pubescens</i>	C, T, o	+	+	+
4	<i>B. pubescens</i>	C, T, p	+	+	+
5	<i>B. pubescens</i>	C, T, p	+	-	+
6	<i>B. pubescens</i>	C, T, d	-	+	+
7	<i>B. pubescens</i>	C, T, p	-	+	na
8	<i>B. pubescens</i>	C, T, p	+	+	na
9	<i>B. pubescens</i>	C, T, p	-	+	+
10	<i>B. pubescens</i>	o	+	+	na
11	<i>B. pubescens</i>	L	-	+	+
12	<i>B. pubescens</i>	C, T, p	+	+	na
13	<i>B. pubescens</i>	C, T, p	-	+	+
14	<i>B. pubescens</i>	C, T, p	+	+	na
15	<i>B. pubescens</i>	C, T, p	+	+	na
16	<i>B. pubescens</i>	C, T, o	-	+	na
17	<i>B. pubescens</i>	C, T, p	+	+	na
18	<i>B. pendula</i>	C, T, p	-	-	na
19	<i>B. pubescens</i>	atypical, p	-	-	na
20	<i>B. pubescens</i>	atypical, p	-	-	na

^{a)} C=clear; T=typical; L=leaves partly fallen; p=symptoms present at time of sampling; d=symptoms decreasing at time of sampling; o=symptoms over at time of sampling

^{b)} +=specific fragment amplification in IC-RT-PCR, -=no amplification, na= sample not available

1997, Osman and Rowhani 2006), may be highly concentrated in senescent tissue.

Electron microscopy revealed isometric virus particles with an estimated diameter of 28–30 nm in leaf samples, originating from *B. pubescens* tree nos. 15 and 17. This is in accordance with the reported morphology of *Cherry leaf roll virus* (Jones 1985), although other viruses as for instance ApMV are known to infect *Betula* species (Grüntzig et al. 1996) exhibiting similar isometric particle sizes. On the contrary no other virus-like structures with different morphologies were visible, so at least filamentous viruses like TRV are unlikely to participate as causal agents of the disease.

This is the first time that *Cherry leaf roll virus* has been confirmed by molecular methods to occur in several *B. pubescens* trees in Rovaniemi and based on symptoms in many birch populations all over Finland. In a previous report on

viruses and mycoplasmas in forest trees in Finland, Bremer et al. (1991) suspected two silver birches (*B. pendula*) at two separate locations in southern Finland to have been CLRV positive by symptom evaluation. In the first case, leaves of the silver birch had pale green lesions and ring spots, which did not appear in recent samples on any birch species. According to Bremer et al. (1991), it was possible to transmit the agent and cause similar symptoms by grafting an infested shoot to a healthy seed-borne silver birch sapling. In the other case, another silver birch with similar symptoms was analysed serologically by J.I. Cooper, who, by personal communication, confirmed that the birch exhibited CLRV (Bremer et al. 1991).

Although Hamacher et al. (1987) has mentioned CLRV-infected *B. pubescens* occurring in Europe before, our positive findings are the first ones on this *Betula* species in Finland. Referring to the visual observation, it seems that most of the trees with CLRV-type symptoms in Finland have been *B. pubescens*, at least so far.

In our countrywide records during the summer of 2006, symptoms on birch appeared very distinct. Although trees were sampled in September, we were able to confirm CLRV infection in birches in Rovaniemi, assuming that this virus is widely spread in Fennoscandia. CLRV is transmitted via pollen and seed in nature (Massalski and Cooper 1984) allowing effective intra-specific dispersal of the virus. While vertical transmission of CLRV by pollen and seed is effective, horizontal transmission by pollen originating from infected birch trees seems to be limited (Cooper et al. 1984). It was also reported by Cooper (1976) that germination rates of birch seedlings from infected mother plants were reduced in comparison with healthy trees, which suggests that vertical transmission is not alone responsible of natural CLRV dissemination and other routes of virus dispersal with epidemiological relevance may exist. The rate at which symptoms have spread in the last few years indicates CLRV to be able to become a serious problem in northern birch forests and for the local forest industry. Therefore rapid efforts are needed to conduct detailed studies on distribution, ways of spread as well as biological and economic importance of CLRV among birch species and possible other hosts. It has to be revealed,

why CLRV is spreading so fast and widely and whether increasing summer temperatures may be responsible for a higher susceptibility to CLRV. The means of viral spread is of special interest because CLRV-typical symptoms were found in a few years time in a large geographical area, in several birch species and on very distinct site types far from each other.

Acknowledgements

The work was supported by Deutsche Forschungsgemeinschaft (DFG, grant no. Bu 890/8-1).

References

- Bandte, M. & Büttner, C. 2001. A review of an important virus of deciduous trees – cherry leaf roll virus: occurrence, transmission and diagnosis. *Pflanzenschutzberichte* 59: 1–19.
- Bremer, K., Lehto, K. & Kurkela, T. 1991. Metsäpuuiden virus- ja mykoplasmatauteja. Diseases caused by viruses and mycoplasmas in forest trees. *Metsäntutkimuslaitoksen tiedonantoja* 382. 15 p.
- Cooper, J.I. 1976. The possible epidemiological significance of pollen and seed transmission in the cherry leaf roll virus/*Betula* spp., complex. *Mitteilungen der Biologischen Bundesanstalt für Land- und Forstwirtschaft, Berlin-Dahlem*, 170: 17–27.
- & Atkinson, M.A. 1975. Cherry leaf roll virus, a cause of chlorotic symptoms in *Betula* spp. in the United Kingdom. *Forestry* 48: 193–203.
- , Massalski, P.R. & Edward, M.L. 1984. Cherry leaf roll virus in the female gametophyte and its relevance to vertical virus transmission. *Annals of Applied Biology* 105: 55–64.
- Demeke, T. & Adams, R.P. 1992. The effects of plant polysaccharides and buffer additives on PCR. *Biotechniques* 12: 332–334.
- Grüntzig, M., Fuchs, E. & Hentsch, T. 1996. Occurrence and serological detection of cherry leaf roll nepovirus (CLRV) and apple mosaic ilarvirus (ApMV) in *Betula* spp. *Journal of Plant Diseases and Protection* 103: 571–581.
- Hamacher, J., Kreutz, G. & Jacob, S. 1987. Histologische Untersuchungen an Kirschenblattrollvirus

- infiizierten Birken. Mededelingen van de Faculteit Landbouwwetenschappen Rijksuniversiteit Gent 52(3a): 1049–1057.
- Harlow, E. & Lane, D. 1988. Antibodies: a laboratory manual. Cold Spring Harbour, N.Y. USA
- Jones, A.T. 1985. CMI/AAB Descriptions of plant viruses no. 306: 6.
- MacKenzie, D.J., Mclean, M.A., Mukerji, S. & Green, M. 1997. Improved RNA extraction from woody plants for the detection of viral pathogens by reverse transcription-polymerase chain reaction. Plant Disease 81: 222–226.
- Massalski P.R. & Cooper J.I. 1984. The location of virus-like particles in the male gametophyte of birch, walnut and cherry naturally infected with cherry leaf roll virus and its relevance to vertical transmission of the virus. Plant Pathology 33: 255–262.
- Monthly weather report. 2006. Finnish Meteorological Institute, Helsinki.
- Osman, F. & Rohwani, A. 2006. Application of a spotting sample preparation technique for the detection of pathogens in woody plants by RT-PCR and real-time PCR (TaqMan). Journal of Virological Methods 133: 130–136.
- Rebenstorf, K., Candresse, T., Dulucq, M.J., Büttner, C. & Obermeier, C. 2006. Host species dependent population structure of a pollen-borne plant virus, Cherry leaf roll virus (CLRV). Journal of Virology 80: 2453–2462.
- Schmelzer, K. 1972. Das Kirschenblattrollvirus aus der Birke (*Betula pendula* Roth.). Zentralblatt für Bakteriologie, Parasitenkunde, Infektionskrankheiten und Hygiene, II. Abteilung 127: 10–12.
- Werner, R., Mühlbach, H.-P. & Büttner, C. 1997. Detection of cherry leaf roll nepovirus (CLRV) in birch, beech and petunia by immunocapture RT-PCR using a conserved primer pair. European Journal of Forest Pathology 27: 309–318.

Total of 17 references